

Human Peripheral Blood B Lymphocytes, #IM-9

SKU: CDS10517 | Category: Cell Lines — Other Established Cell Lines

Validation & Biomarker Data Guide · Lineage classification: B lymphocyte

1. Product Overview

Human Peripheral Blood B Lymphocytes, #IM-9 is an established cell line for controlled in vitro research. It can support cell biology, growth and viability studies, signaling and pathway research, drug-response testing, screening and assay development under validated culture conditions. The model provides a reproducible cellular system for basic and translational research. For research use only.

Primary tissue / source: Peripheral Blood B Lymphocytes | Growth type: Adherent Cells | Tumorigenic status: Normal Cells | Pack size: 1 x 1 million cells

2. Typical Applications

- 1. Cancer Biology Research
- Mechanistic Studies
- Gene Function Studies
- Signal Transduction
- 2. Drug Discovery and Development
- Screening of Anticancer Compounds
- IC50/EC50 Studies
- Mechanism of Action
- Combination Therapies
- 3. Assay Development
- ELISA and Immunoassays
- Flow Cytometry Assays

3. Identity & Phenotypic Marker Panel

A representative identity panel for b lymphocyte products is shown below. Actual per-lot Certificates of Analysis should be consulted for verified identity data.

Marker	Expected Result	Range
CD19	Positive	80–98% positive

Marker	Expected Result	Range
CD20	Positive	75–97% positive
HLA-DR	Positive	70–95% positive
CD3	Negative / low	≤5% (background)
CD14	Negative / low	≤4% (background)




Figure 1. Marker expression profile (mean ± SD, n = 3 replicates).

4. Growth Kinetics & Viability

Growth and viability trends over 120 hours in standard culture conditions.




Figure 2. Growth curve (viable cell density, left axis) and viability trend (right axis).

5. Functional Validation Assay

B-cell activation assay: Stimulation with anti-IgM/CpG or CD40L to assess proliferation and immunoglobulin secretion (IgM/IgG) by ELISA.




Figure 3. Functional assay readout — ~3.9× induction over baseline.

6. QC Acceptance Criteria

Acceptance Criterion	Result
≥80% CD19+ by flow cytometry	PASS
Immunoglobulin secretion confirmed by ELISA	PASS
Negative for mycoplasma by PCR	PASS
Apparent population doubling time	≈ 44 hours
Functional assay outcome	~3.9× induction over baseline

7. Handling & Culture Notes

- Refer to the product-specific Certificate of Analysis and datasheet for validated culture medium, seeding density, and passaging recommendations.
- As with any primary cell or cell line, confirm mycoplasma-negative status, expected morphology by phase-contrast microscopy, and viability by trypan blue or equivalent dye-exclusion assay at each passage before use in downstream assays.