

Primary Rabbit Coronary Artery Endothelial Cells (CAEC)

SKU: CDS11244 | Category: Primary Cells — Rabbit

Validation & Biomarker Data Guide · Lineage classification: Endothelial cell

1. Product Overview

Primary Rabbit Coronary Artery Endothelial Cells (CAEC) is a primary cell product from rabbit Coronary artery for physiologically relevant in vitro studies. The product is particularly relevant to Cardiac cell series and Endothelial cells research. It can support research in cell physiology, signaling, metabolism, toxicology, drug response and cell-based assay development where a primary-cell model is preferred. Culture and passage conditions should follow the product-specific protocol. For research use only.

Primary tissue / source: Coronary artery | Growth type: Adherent Cells | Tumorigenic status: Normal Cells
| Pack size: 1 x 0.5 million cells

2. Typical Applications

- Cardiac cell series
- Endothelial cells
- Circulatory system

3. Identity & Phenotypic Marker Panel

A representative identity panel for endothelial cell products is shown below. Actual per-lot Certificates of Analysis should be consulted for verified identity data.

Marker	Expected Result	Range
CD31 (PECAM-1)	Positive	85–99% positive
VE-cadherin	Positive	75–97% positive
vWF	Positive	70–95% positive
Pan-cytokeratin	Negative / low	≤5% (background)
α-SMA	Negative / low	≤10% (background)




Figure 1. Marker expression profile (mean ± SD, n = 3 replicates).

4. Growth Kinetics & Viability

Growth and viability trends over 120 hours in standard culture conditions.




Figure 2. Growth curve (viable cell density, left axis) and viability trend (right axis).

5. Functional Validation Assay

Tube formation (angiogenesis) assay: Matrigel tube-formation assay quantifying network/branch-point formation to confirm angiogenic functional competence; Ac-LDL uptake as secondary marker.




Figure 3. Functional assay readout — $\sim 5.2\times$ induction over baseline.

6. QC Acceptance Criteria

Acceptance Criterion	Result
$\geq 85\%$ CD31+ by flow cytometry/ICC	PASS
Positive tube formation on Matrigel within 6-18 h	PASS
Apparent population doubling time	≈ 29 hours
Functional assay outcome	$\sim 5.2\times$ induction over baseline

7. Handling & Culture Notes

- Refer to the product-specific Certificate of Analysis and datasheet for validated culture medium, seeding density, and passaging recommendations.
- As with any primary cell or cell line, confirm mycoplasma-negative status, expected morphology by phase-contrast microscopy, and viability by trypan blue or equivalent dye-exclusion assay at each passage before use in downstream assays.