

Primary Canine Hepatocyte (Hep)

SKU: CDS11982 | Category: Primary Cells — Dog

Validation & Biomarker Data Guide • Lineage classification: Hepatocyte

1. Product Overview

Primary Canine Hepatocyte (Hep) is a primary cell product from canine Liver for physiologically relevant in vitro studies. The product is particularly relevant to Hepatology and Hepatic cell series research. It can support research in cell physiology, signaling, metabolism, toxicology, drug response and cell-based assay development where a primary-cell model is preferred. Culture and passage conditions should follow the product-specific protocol. For research use only.

Primary tissue / source: Liver | Growth type: Adherent Cells | Tumorigenic status: Normal Cells | Pack size: 1 x 0.5 million cells

2. Typical Applications

- Hepatology
- Hepatic cell series
- Hepatocytes
- Parenchymal cells
- Hemolymphatic system

3. Identity & Phenotypic Marker Panel

A representative identity panel for hepatocyte products is shown below. Actual per-lot Certificates of Analysis should be consulted for verified identity data.

Marker	Expected Result	Range
Albumin	Positive	70–96% positive
HNF4-α	Positive	65–93% positive
CYP3A4 activity	Positive	60–90% positive
CD45	Negative / low	≤3% (background)
Vimentin	Negative / low	≤20% (background)




Figure 1. Marker expression profile (mean $\pm$ SD, n = 3 replicates).

4. Growth Kinetics & Viability

Growth and viability trends over 120 hours in standard culture conditions.




Figure 2. Growth curve (viable cell density, left axis) and viability trend (right axis).

5. Functional Validation Assay

CYP450 metabolic activity assay: Luminescent or fluorometric CYP3A4/CYP1A2 activity assay following inducer exposure (e.g. rifampicin); urea and albumin secretion quantified by ELISA as secondary liver-function markers.




Figure 3. Functional assay readout — ~3.3× induction over baseline.

6. QC Acceptance Criteria

Acceptance Criterion	Result
Albumin secretion above lineage reference range (ELISA)	PASS
Detectable inducible CYP3A4 activity	PASS
Urea synthesis confirmed by colorimetric assay	PASS
Apparent population doubling time	≈ 51 hours
Functional assay outcome	~3.3× induction over baseline

7. Handling & Culture Notes

- Refer to the product-specific Certificate of Analysis and datasheet for validated culture medium, seeding density, and passaging recommendations.
- As with any primary cell or cell line, confirm mycoplasma-negative status, expected morphology by phase-contrast microscopy, and viability by trypan blue or equivalent dye-exclusion assay at each passage before use in downstream assays.