


Human K562 Doxorubicin-Resistant Cell Line, #K562/ADR

Human K562 Doxorubicin-Resistant Cell Line, #K562/ADR is an engineered or reporter cell line for controlled in vitro studies. The resistant phenotype provides a basis for investigating how cancer cells adapt to treatment pressure.

Applications include comparative drug-response testing and investigation of resistance mechanisms using appropriate parental controls.

CTDS SKU	CDS10119
Pack size	1 x 1 million cells
Product category	Engineered Cell Lines
Product subcategory	Drug-Resistant Cell Lines

PRODUCT IMAGE



PRODUCT BACKGROUND

Human K562 Doxorubicin-Resistant Cell Line, #K562/ADR is an engineered or reporter cell line for controlled in vitro studies. It is suitable for cell-based assay development, pathway analysis, target-response studies, screening, pharmacology and comparative functional testing. Engineered features stated in the product name can support quantitative or selectable experimental readouts. For research use only.

APPLICATIONS

1. Cancer Biology Research
 - Mechanistic Studies
 - Gene Function Studies
 - Signal Transduction
2. Drug Discovery and Development
 - Screening of Anticancer Compounds
 - IC50/EC50 Studies
 - Mechanism of Action
 - Combination Therapies
3. Assay Development
 - ELISA and Immunoassays
 - Flow Cytometry Assays
 - Reporter Gene Assays
4. Omics and Biomarker Research
 - Transcriptomics / Genomics
 - Proteomics
 - Metabolomics
5. Tumor Heterogeneity and Resistance Mechanisms
 - Clonal Selection

Acquired Resistance Models

6. Personalized Medicine and Precision Oncology

Patient-Derived Cell Lines

Companion Diagnostics

7. Immuno-oncology

Co-culture with Immune Cells

Checkpoint Inhibitor Testing

Antibody-Dependent Cytotoxicity (ADCC)

8. Vaccine Development

Neoantigen Discovery

Antigen Presentation Studies

BIOLOGICAL CHARACTERISTICS

Alternate name 1	CVCL_1938 Cells
Alternate name 2	K562/DOX, K562/Adriamycin Cells
Cell growth type	Suspension Cells
Biosafety level	BSL-2
Tumorigenicity / cell status	Tumor Cells
Target / construct	Drug-resistant derivative: Human K562 Doxorubicin-Resistant Cell Line

CULTURE REQUIREMENTS

Culture medium	RPMI1640 + 10% FBS + 1% Pen/Strep + 1 ug/ml ADR
Freezing medium	55% Complete Culture Medium + 40% FBS + 5% DMSO

RECEIPT AND CRYOPRESERVATION

Upon Receipt of Cells

1 mL cryovials are packed in dry ice for transportation. After receipt, clean the cryovial with 70% IPA to disinfect. Stored in a -80°C refrigerator overnight and then transfer to liquid nitrogen or revive them directly as per your use.

Note If the dry ice is found to have evaporated or the cryovial cap is detached/ damaged or the cells seems damaged, please contact us immediately.

CULTURE AND RECOVERY INSTRUCTIONS

Recovery of Frozen Cells:

For Suspension Cell passaging, the following methods can be used as reference:

- 1) Shake the cryotube containing 1mL of cell suspension quickly within seconds in a 37°C water bath to thaw.
- 2) Add it to a centrifuge tube containing 4-6mL of complete medium and mix well. Centrifuge at 1000rpm for 3-5mins.
- 3) Discard the supernatant, and resuspend the cells in complete medium.
- 4) Then add the cell suspension to a culture flask (or dish) containing 6-8ml of complete medium and culture overnight at 37°C.
- 5) Observe the cell growth and cell density under a microscope the next day.
- 6) Cell subculture: If the cell density reaches 80%-90%, subculture can be performed.

For Adherent Cell passaging, the following methods can be used as reference:

- 1) Shake the cryotube containing 1mL of cell suspension quickly within seconds in a 37°C water bath to thaw.
- 2) Add it to a centrifuge tube containing 4-6mL of complete medium and mix well. Centrifuge at 1000rpm for 3-5mins.
- 3) Discard the supernatant, and rinse the cells 1-2 times with PBS without calcium and magnesium ions.
- 4) Add 0.25% (w/v) trypsin-0.53 mM EDTA to the culture flask (1-2 mL for T25 flask, 2-3 mL for T75 flask),
- 5) Place in a 37°C incubator for digestion for 1-2 minutes (difficult-to-digest cells can be properly prolonged for digestion time),
- 6) Observe the cell digestion under a microscope. If most of the cells become round and fall off, quickly take it back to the operating table, tap the culture flask a few times, and add 3-4 ml of culture medium containing 10% FBS to terminate digestion.

- 7) Gently mix and aspirate, centrifuge at 1000rpm for 3-5min, discard the supernatant. Add 1-2mL culture medium and mix evenly.
- 8) Divide the cell suspension into new T25 bottles at a ratio of 1:2.
- 9) Add 6-8ml of new complete culture medium prepared according to the instructions to maintain cell growth vitality, and perform subsequent passages at a ratio of 1:2~1:5 as per actual conditions.

CHARACTERIZATION AND QUALITY INFORMATION

Sterility	Bacteria: Negative
	Yeast: Negative
	Mycoplasma: Negative
Pathogen testing	HIV: Negative
	Hepatitis B: Negative
Markers / assay readout	T-lineage panel: CD45+, surface/cytoplasmic CD3+, CD4 and CD8, CD7+, CD5+; add viability dye. Confirm line-specific Phenotype.
Assay / validation method	Comparative drug-response / resistance assay; Flow cytometry marker characterization
Authentication / QC evidence	T-lineage panel: CD45+, surface/cytoplasmic CD3+, CD4 and CD8, CD7+, CD5+; add viability dye. Confirm line-specific Phenotype.; Sterility: Bacteria: Negative
	Yeast: Negative
	Mycoplasma: Negative; Pathogen testing: HIV: Negative Hepatitis B: Negative

Catalogue confirmation note: Product-specific confirmation required: list only markers actually tested. Current text describes a suggested panel.

HANDLING AND STORAGE

Shelf life	12 months at time of shipping
Shipping	Shipped on Dry Ice at -20°C
Storage	Store at -176°C.
Intended use	For Research Use Only. Not for diagnostics or human use.

PRODUCT DISCLAIMER

The data indicated herein are as indicated and validated in our laboratory. As these cell lines are of human origin, they carry a risk of latent human viruses (e.g., Epstein-Barr virus, hepatitis B). Therefore they should be handled accordingly with precaution. These reagents are for research use only and not for in-vitro diagnostics or human use.

ADDITIONAL CLASSIFICATION

Product grouping	Engineered Cell Lines
Product subgroup	Human

Not a lot-specific certificate of analysis.