

## LS1034 Human Colon Carcinoma Cell Line

LS1034 Human Colon Carcinoma Cell Line is an established tumor-derived cell line for in vitro oncology research. Intestinal-cancer models support research into tumor-cell renewal, survival and responses to treatment. Applications include studies of proliferation, survival signaling, cytotoxicity and comparative responses to experimental treatments.

|                     |                     |
|---------------------|---------------------|
| CTDS SKU            | CDS1525             |
| Pack size           | 1 x 1 million cells |
| Product category    | Cell Lines          |
| Product subcategory | Cancer Cell Lines   |

### PRODUCT BACKGROUND

LS1034 Human Colon Carcinoma Cell Line is an established tumor-derived cell line for in vitro oncology research. It is suitable for studies of tumor cell biology, proliferation and viability, signaling, biomarkers, drug response, cytotoxicity and cell-based assay development. The model can be incorporated into screening and mechanistic workflows under validated culture conditions. For research use only.

### BIOLOGICAL CHARACTERISTICS

|                              |                |
|------------------------------|----------------|
| Host cell                    | LS1034         |
| Alternate name 1             | Cell Line      |
| Cell growth type             | Adherent Cells |
| Biosafety level              | BSL-2          |
| Tumorigenicity / cell status | Tumor Cells    |

### CULTURE REQUIREMENTS

|                 |                                                                                                                                             |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| Culture medium  | DMEM + 10% FBS as a general starting formulation; use the product-specific CTDS complete medium where a dedicated formulation is specified. |
| Freezing medium | 90% FBS + 10% DMSO                                                                                                                          |

### RECEIPT AND CRYOPRESERVATION

#### Upon Receipt of Cells

1 mL cryovials are packed in dry ice for transportation. After receipt, clean the cryovial with 70% IPA to disinfect. Stored in a -80°C refrigerator overnight and then transfer to liquid nitrogen or revive them directly as per your use.

Note If the dry ice is found to have evaporated or the cryovial cap is detached/ damaged or the cells seems damaged, please contact us immediately.

### CULTURE AND RECOVERY INSTRUCTIONS

#### Recovery of Frozen Cells:

For Suspension Cell passaging, the following methods can be used as reference:

- 1) Shake the cryotube containing 1mL of cell suspension quickly within seconds in a 37°C water bath to thaw.
- 2) Add it to a centrifuge tube containing 4-6mL of complete medium and mix well. Centrifuge at 1000rpm for 3-5mins.
- 3) Discard the supernatant, and resuspend the cells in complete medium.
- 4) Then add the cell suspension to a culture flask (or dish) containing 6-8ml of complete medium and culture overnight at 37°C.
- 5) Observe the cell growth and cell density under a microscope the next day.
- 6) Cell subculture: If the cell density reaches 80%-90%, subculture can be performed.

For Adherent Cell passaging, the following methods can be used as reference:

- 1) Shake the cryotube containing 1mL of cell suspension quickly within seconds in a 37°C water bath to thaw.
- 2) Add it to a centrifuge tube containing 4-6mL of complete medium and mix well. Centrifuge at 1000rpm for 3-5mins.
- 3) Discard the supernatant, and rinse the cells 1-2 times with PBS without calcium and magnesium ions.

- 4) Add 0.25% (w/v) trypsin-0.53 mM EDTA to the culture flask (1-2 mL for T25 flask, 2-3 mL for T75 flask),
- 5) Place in a 37°C incubator for digestion for 1-2 minutes (difficult-to-digest cells can be properly prolonged for digestion time),
- 6) Observe the cell digestion under a microscope. If most of the cells become round and fall off, quickly take it back to the operating table, tap the culture flask a few times, and add 3-4 ml of culture medium containing 10% FBS to terminate digestion.
- 7) Gently mix and aspirate, centrifuge at 1000rpm for 3-5min, discard the supernatant. Add 1-2mL culture medium and mix evenly.
- 8) Divide the cell suspension into new T25 bottles at a ratio of 1:2.
- 9) Add 6-8ml of new complete culture medium prepared according to the instructions to maintain cell growth vitality, and perform subsequent passages at a ratio of 1:2~1:5 as per actual conditions.

## CHARACTERIZATION AND QUALITY INFORMATION

|                              |                                                                             |
|------------------------------|-----------------------------------------------------------------------------|
| Sterility                    | Bacteria: Negative                                                          |
|                              | Yeast: Negative                                                             |
|                              | Mycoplasma: Negative                                                        |
| Pathogen testing             | HIV: Negative                                                               |
|                              | Hepatitis B: Negative                                                       |
| Authentication / QC evidence | Sterility: Bacteria: Negative                                               |
|                              | Yeast: Negative                                                             |
|                              | Mycoplasma: Negative; Pathogen testing: HIV: Negative Hepatitis B: Negative |

## HANDLING AND STORAGE

|              |                                                          |
|--------------|----------------------------------------------------------|
| Shelf life   | 12 months at time of shipping                            |
| Shipping     | Shipped on Dry Ice at -20°C                              |
| Storage      | Store at -176°C.                                         |
| Intended use | For Research Use Only. Not for diagnostics or human use. |

## PRODUCT DISCLAIMER

The data indicated herein are as indicated and validated in our laboratory. As these cell lines are of human/animal origin, they carry a risk of latent human/animal viruses (e.g., Epstein-Barr virus, hepatitis B). Therefore they should be handled accordingly with precaution. These reagents are for research use only and not for in-vitro diagnostics or human use.

## ADDITIONAL CLASSIFICATION

|                  |                   |
|------------------|-------------------|
| Product grouping | Cancer Cell Lines |
|------------------|-------------------|

Not a lot-specific certificate of analysis.